



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*On the Error of a received Principle of Analysis, respecting Functions which vanish with their Variables.* By WILLIAM R. HAMILTON, *Royal Astronomer of Ireland, &c.*

Read January 25, 1830.

---

It appears to be a received principle of analysis, that if a real function of a positive variable ( $x$ ) approaches to zero with the variable, and vanishes along with it, then that function can be developed in a real series of the form

$$Ax^{\alpha} + Bx^{\beta} + Cx^{\gamma} + \&c.,$$

the exponents  $\alpha, \beta, \gamma, \dots$  being constant and positive, and the coefficients  $A, B, C, \dots$  being constant, and all these constant exponents and coefficients being finite and different from zero. This principle has been made the foundation of important theories, and has not ever, so far as I know, been questioned; but I believe that the following example of exception, which it would be easy to put in a more general form, will sufficiently prove it to be erroneous; since if the principle be true, it is by its nature universal.

The real function  $e^{-x^{-2}}$ , in which  $e$  is the base of the neperian logarithms, approaches to zero along with  $x$  and vanishes along with it. Yet if we could develop this function in a series of the kind described, we should have

$$x^{-\alpha} e^{-x^{-2}} = A + Bx^{\beta-\alpha} + Cx^{\gamma-\alpha} + \&c.$$

in which we might suppose  $\alpha$  the least exponent; and then, while  $x$  approached to 0, the second member would tend to the limit  $A$ , which by hypothesis is different from 0; and yet, from the nature of exponential functions, the limit of the first member is zero.

We conclude, therefore, that the function  $e^{-x^{-\alpha}}$  cannot be developed in a series of the kind assumed, although it vanishes with its variable; and consequently that, if we only know this property of a function, that it vanishes when its variable vanishes, we cannot correctly assume that it may be developed in such a series.

If any doubt should be felt respecting the truth of the remark, that the function  $x^{-\alpha} e^{-x^{-\alpha}}$  tends to zero along with  $x$ , when  $\alpha$  is any positive constant, this doubt will be removed by observing that the function  $x^{\alpha} e^{x^{-\alpha}}$ , which is the reciprocal of the former, increases without limit while  $x$  decreases to zero. For we may develop this latter function, by the known theorems, in the essentially converging series,

$$x^{\alpha} e^{x^{-\alpha}} = y^{-\alpha} e^{y^{\alpha}} = y^{-\alpha} + y^{2-\alpha} + \frac{y^{4-\alpha}}{2} + \frac{y^{6-\alpha}}{2 \cdot 3} + \frac{y^{8-\alpha}}{2 \cdot 3 \cdot 4} + \&c.,$$

$y$  being the reciprocal of  $x$ ; and while  $y$  tends to  $+\infty$ , the terms of this series remain all positive, and all after a certain constant number increase indefinitely.